

ISSN 2221-6421 (Print), ISSN 2223-2567 (Online)

Coden: PJSIC6 61(2) 59-120 (2018)

# **Pakistan Journal of Scientific and Industrial Research**

**Series B: Biological Sciences**

**Vol. 61, No.2, May-August, 2018**



Published by  
Scientific Information Centre  
Pakistan Council of Scientific and Industrial Research  
Karachi, Pakistan

# Pakistan Journal of Scientific and Industrial Research

## Series B: Biological Sciences

### EDITORIAL BOARD

**Dr. Shahzad Alam**

Chief Editor

**Shagufta Yasmin Iqbal**

Executive Editor

### MEMBERS

**Prof. Z. Adamek**

Faculty of Agriculture  
Univ. South Bohemia,  
Czech Republic

**Dr. T. A. Ajith**

Amala Institute of Medical Science  
Kerala, India

**Dr. S. A. Barlas**

Environ. Protection Division  
Penticton, BC V0H 1Z4,  
Canada

**Dr. C. M. A. Caipang**

Bio Vivo Technologies As, Norway

**Dr. D. Das**

C. B. Fleet Company, Inc., VA,  
USA

**Prof. M. J. Jaskani**

Plant Tissue Culture Cell  
Univ. of Agric., Faisalabad, Pakistan

**Dr. A. Khanum**

Dept. Biochemistry  
Pir Mehr Ali Shah Agric. University  
Rawalpindi, Pakistan

**Dr. S. Mathews**

Pharmacology, The University of  
Sydney, NSW, Australia

**Prof. E. Miraldi**

Pharmaceutical Biology Section  
University of Siena, Siena, Italy

**Dr. G. Müller**

Sanofi-Aventis Pharma  
Frankfurt, Germany

**Dr. S. K. Rastogi**

Dept. of Chem. & Biochemistry  
Texas State University, USA

**Dr. J. S. Sangha**

Dalhousie University Truro,  
NS B2N 5E3, Canada

**Dr. H. Shimoda**

Oryza Oil & Fat Chemical Co. Ltd.,  
Aichi, Japan

**Prof. Dr. T. Toyosaki**

Dept. of Foods and Nutrition  
Fukuoka, Japan

**Dr. V. Zambare**

Centre for Bioprocessing  
Research and Development,  
South Dakota, USA

**Editors:** Shahida Begum Sajid Ali Seema Iqbal

Pakistan Journal of Scientific and Industrial Research started in 1958, has been bifurcated in 2011 into:

**Series A: Physical Sciences** [ISSN 2221-6413 (Print); ISSN 2223-2559 (online)] (appearing as issues of January-February, May-June and September-October) and

**Series B: Biological Sciences** [ISSN 2221-6421 (Print); ISSN 2223-2567 (online)] (appearing as issues of March-April, July-August and November-December).

Each Series will appear three times in a year.

**This Journal is indexed/abstracted** in Biological Abstracts and Biological Abstracts Reports, Chemical Abstracts, Geo Abstracts, CAB International, BioSciences Information Service, Zoological Record, BIOSIS, NISC, NSDP, Current Contents, CCAB, Rapra Polymer Database, Reviews and Meetings and their CD-ROM counterparts etc.

**Subscription rates (including handling and Air Mail postage):** *Local:* Rs. 2500 per volume, single issue Rs. 425; *Foreign:* US\$ 450 per volume, single issue US\$ 75.

**Electronic format** of this journal is available with: ProQuest Information and Learning, 789 E. Eisenhower Parkway, P.O. Box 1346, Ann Arbor, MI 48106-1346, U.S.A.; Fax.No.+1.734.997.4268; <http://www.proquest.com>.

**Photocopies of back issues** can be obtained through submission of complete reference to the Executive Editor against the payment of Rs. 25 per page per copy (by Registered Mail) and Rs. 115 per copy (by Courier Service), within Pakistan; US\$ 10 per page per copy (by Registered Mail) and US\$25 per page per copy (by Courier Service), for all other countries.

**Copyrights** of this Journal are reserved; however, limited permission is granted to researchers for making references, and libraries/agencies for abstracting and indexing purposes according to the international practice.

**Printed and Published by:** PCSIR Scientific Information Centre, PCSIR Laboratories Campus, Shahrah-e-Dr. Salimuzzaman Siddiqui, Karachi-75280, Pakistan.

### Editorial Address

#### Executive Editor

Pakistan Journal of Scientific and Industrial Research, PCSIR Scientific Information Centre,  
PCSIR Laboratories Campus, Shahrah-e-Dr. Salimuzzaman Siddiqui, Karachi-75280, Pakistan

Tel: 92-21-99261914-99261916, 99261949-99261917; Fax: 92-21-99261913; Web: <http://www.pjsir.org>, E-mail: [info@pjsir.org](mailto:info@pjsir.org)

## AIMS & SCOPE

Pakistan Journal of Scientific and Industrial Research ( PJSIR ) was started in 1958 to disseminate research results based on utilization of locally available raw materials leading to production of goods to cater to the national requirements and to promote S&T in the country. Over the past 60 years, the journal conveys high quality original research results in both basic and applied research in Pakistan. A great number of major achievements in Pakistan were first disseminated to the outside world through PJSIR.

It is a peer reviewed journal and published in both print and electronic form. Original research articles, review articles and short communications from varied key scientific disciplines are accepted however, papers of Pure Mathematics, Computer Sciences, Engineering and Medical Sciences are not entertained.

From 54<sup>th</sup> Volume in 2011, it has been bifurcated into Series A: Physical Sciences & Series B: Biological Sciences. Each series appears three times in a year as follows:

Physical Sciences in January-February, May-June and September-October issues. It includes research related to Natural Sciences, Organic Chemistry, Inorganic Chemistry, Industrial Chemistry, Physical Chemistry, Environmental Sciences, Geology, Physics, Polymer Sciences and Technology.

Biological Sciences in March-April, July-August and November-December issues. Papers included in this series are from Agriculture, Agronomy, Botany, Biochemistry, Biotechnology, Food Sciences, Genetic Engineering, Pharmaceutical Sciences, Microbiology, Marine Sciences, Soil Sciences, Tissue Culture, Zoology and Technology.

Due to many global issues, we are encouraging contributions from scientists and researchers from all across the globe with the sole purpose of serving scientific community worldwide on the whole and particularly for our region and third world countries.

---

**Pakistan Journal of Scientific and Industrial Research**  
**Series B: Biological Sciences**  
**Vol. 61, No. 2, May-August, 2018**

---

**Contents**

<b>Sorption-Desorption Characteristics of Benzimidazole Based Fungicide Benomyl on Physicochemical Properties of Selected Pakistani Soils and their Minerals</b> Khuram Shahzad Ahmad	59
<b>Response of Wheat to Gypsum Application at Farmers Fields in Rainfed Pothwar</b> Adnan Umair, Muhammad Nadeem Iqbal, Kashif Bashir, Khadim Hussain, Waqas Naseem, Muhammad Rafique Sajjad, Tariq Mehmood and Ghulam Qadir	68
<b>Impact of Different Shapes of Pitchers on Water Saving and Water Use Efficiency of Ridge-gourd in Semi Arid Region of Pakistan</b> Ashifa Soomro, Kamran Baksh Soomro, Azhar Ali Laghari and Samina Siddiqui	72
<b>Development and Characterisation of Fig (<i>Ficus carica</i>) Leaves Tea with Special Reference to Hypoglycemic Effect</b> Maham Gillani, Hira Iftikhar, Imran Pasha and Asma Lodhi	78
<b>Optimisation of Production Condition of Alkaline Protease Using Indigenous <i>Bacillus subtilis</i> from Agricultural Soil</b> Barkat Ali Solangi, Hafiz Rub Nawaz, Uzma Nadeem, Beena Zehra and Agha Asad Noor	84
<b>Deforestation Drivers in Tehsil Barawal, Dir Upper, Pakistan</b> Anwar Sajjad, Syed Adnan and Ahmad Hussain	89
<b>Population Density Estimates of Punjab Urial in Scrub Forest</b> Muhammad Irshad Arshad and Anwar Hussain	95
<b>Comparative Toxicity of Spinetoram and Nitenpyram against Earthworm and their Effects on Protein Contents and Cholinesterase Activity</b> Altaf Hussain, Muhammad Farhanullah Khan, Omer Mukhtar, Muhammad Faheem and Umeed Ali	103
<b>Short Communication</b>	
<b>Correlation and Regression Coefficient Estimates between Some Growth Performance Traits of Harnai Sheep</b> Rameez Raja Kaleri, Hubdar Ali Kaleri, Asma Kaleri, Abdul Rashid, Rashid Ali Shah, Gul Muhammad Marri, Rajesh Kumar and Deepesh Kumar	112
<b>Review</b>	
<b>A Review on Rangeland Management in Pakistan, Bottlenecks and Recommendations</b> Muhammad Jamil, Muhammad Mansoor, Fawad Anwar, Sher Muhammad and Aftab Ahmed Awan	115

# **Sorption-Desorption Characteristics of Benzimidazole Based Fungicide Benomyl on Physicochemical Properties of Selected Pakistani Soils and their Minerals**

**Khuram Shahzad Ahmad**

Department of Environmental Sciences, Fatima Jinnah Women University, Rawalpindi-46000, Pakistan

(received April 1, 2015; revised July 4, 2017; accepted July 10, 2017)

---

**Abstract.** A versatile cost-effective fungicide, Benomyl (Methyl{1(butylamino)carbonyl}-1H-benzimidazol-2-ylcarbamate) has been utilised to investigate its sorption behaviour on physicochemical properties of soils, via batch equilibrium method. Linear and Freundlich adsorption isotherms were evaluated in two tested soils having values of slope  $n < 1$  resembling the c-type curve. The distribution co-efficient  $K_{ads}$  for adsorption was 2.93 and 14.35 mL/ $\mu$ g both soils indicating low adsorption of Benomyl overall with relatively greater degree of adsorption on hilly soil and silt loam and lesser adsorption on sandy soil. Multan soils have more sand content (52%) so it has less  $K_d$  value. Desorption studies revealed that the adsorbed fungicides were firmly retained by soil particles and their adsorption was almost irreversible. Furthermore, the results were statistically evaluated through regression analysis and univariate ANOVA, also probability graphs were plotted to ensure the accuracy of the experimental data. The XRD analysis depicted an increase in basal spacing of pesticide in both soils.

**Keywords:** benomyl, sorption, desorption, physicochemical properties, soil organic matter

---

## Response of Wheat to Gypsum Application at Farmers Fields in Rainfed Pothwar

Adnan Umair<sup>a\*</sup>, Muhammad Nadeem Iqbal<sup>b</sup>, Kashif Bashir<sup>a</sup>, Khadim Hussain<sup>c</sup>, Waqas Naseem<sup>d</sup>,  
Muhammad Rafique Sajjad<sup>c</sup>, Tariq Mehmood<sup>d</sup> and Ghulam Qadir<sup>e</sup>

<sup>a</sup>Soil and Water Conservation Research Station, Sohawa, Pakistan

<sup>b</sup>Soil Fertility Research Institute, Lahore, Pakistan

<sup>c</sup>Soil Testing Laboratory, Jhelum, Pakistan

<sup>d</sup>Soil and Water Conservation Research Institute, Chakwal, Pakistan

<sup>e</sup>Soil Salinity Research Institute, Pindi Bhatian, Pakistan

(received February 17, 2016; revised February 20, 2017; accepted February 21, 2017)

---

**Abstract.** In present study, possibility of use and dose of gypsum was explored in various agro-ecologies of Pothwar region, Pakistan. Experiments were conducted at farmers' fields for three years i.e., 2002-2003 to 2004-2005 at three different districts viz. Chakwal, Attock, Jhelum and Rawalpindi. After first rainfall of monsoon in the end of June, deep ploughing was done in the field with mouldboard plough. After that, gypsum was applied @ 1.25, 2.5, 5.0 and 7.5 t/ha to experimental plots. Application of gypsum resulted in increase in grain yield of wheat during three years. Application of gypsum increased grain yield in all the districts. There was 11.9 to 34 percent increase in wheat yield with application of 2.5 t/ha of gypsum. At experimental plots, where grain yields were less than 3.1 t/ha after application of 2.5 t/ha of gypsum, increasing gypsum application beyond 2.5 t/ha increased the yield further. However, at sites, where grain yields were more than 3.4 t/ha with application of gypsum @ 2.5 t/ha, further increase in gypsum dose beyond 2.5 t/ha decreased the yields. On the basis of these experiments, 2.5 t/ha was found the appropriate dose of gypsum for rainfed wheat.

**Keywords:** gypsum, wheat, rainfed, rainfall, Pothwar

---

# Impact of Different Shapes of Pitchers on Water Saving and Water Use Efficiency of Ridge-gourd in Semi Arid Region of Pakistan

Ashifa Soomro<sup>a</sup>, Kamran Baksh Soomro<sup>\*b</sup>, Azhar Ali Laghari<sup>a</sup> and Samina Siddiqui<sup>c</sup>

<sup>a</sup>Department of Land and Water Management, Sindh Agriculture University Tandojam, Sindh, Pakistan

<sup>b</sup>Civil Engineering Discipline, School of Engineering, Monash University, Malaysia

<sup>c</sup>National Centre for Excellence in Geology, University of Peshawar, KPK, Pakistan

(received December 3, 2016; revised December 5, 2017; accepted February 13, 2018)

---

**Abstract.** Pitcher irrigation is an ancient irrigation system, used for cultivation of small areas where water is saline and not fit for surface irrigation. This research aims to investigate the effect on water use efficiency and water saving of ridge gourd by utilization of different shapes of pitchers. Ridge-gourd was planted under four different pitcher shapes including T<sub>1</sub> (conical), T<sub>2</sub> (spherical), T<sub>3</sub> (round) and T<sub>4</sub> (cylindrical) shapes at the experimental field of Sindh Agriculture University Tandojam, Pakistan. Overall sixteen clay pitchers having 1.7, 1.3, 1.3 and 1.08 m<sup>3</sup> volume of water under T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub> respectively were installed in the soil. The irrigation water having electrical conductivity (EC) 1.3 dS/m with pH of 7.8 was used to fill the pitchers. The highest soil EC of 3.32 dS/m at 0-20 cm depth of soil was found under T<sub>4</sub> followed by T<sub>3</sub>, T<sub>2</sub> and T<sub>1</sub>, respectively. Whereas T<sub>4</sub> showed minimum soil EC as 1.30 dS/m at the depth of 40-60 cm. However the pH effect of T<sub>1</sub> was slightly higher as 8.4 at the depth of 40-60 cm in comparison to T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub>, whereas the minimum pH of 8.0 was found at 20-40 cm under T<sub>4</sub>. The maximum vine length of ridge gourd was 211cm under T<sub>4</sub> and the minimum length of vine was 139 cm under T<sub>2</sub>. The highest water use efficiency 8.6 kg/m<sup>3</sup> was obtained under T<sub>4</sub> whereas the highest water saving as 82% was recorded under T<sub>4</sub> followed by T<sub>2</sub>, T<sub>1</sub> and T<sub>3</sub>.

**Keywords:** Pitcher irrigation, Ridge gourd, water use efficiency, water saving, vine length

---

## Development and Characterisation of Fig (*Ficus carica*) Leaves Tea with Special Reference to Hypoglycemic Effect

Maham Gillani<sup>a\*</sup>, Hira Iftikhar<sup>a</sup>, Imran Pasha<sup>b</sup> and Asma Lodhi<sup>a</sup>

<sup>a</sup>Institute of Home Sciences, University of Agriculture, Faisalabad, Pakistan

<sup>b</sup>National Institute of Food Science and Technology, University of Agriculture, Faisalabad, Pakistan

(received December 2, 2016; revised July 26, 2017; accepted August 1, 2017)

---

**Abstract.** The leaves of *Ficus carica* (Fig) possess wide range of biological activities and are used to treat diabetes and its complications. The aim of the study was to evaluate the hyperglycemic and antioxidant effect of tea of *Ficus carica* leaves on normal and hyperglycemic rats. Effect of fig leaves tea was investigated on blood glucose level, daily food intake, daily water intake, body weight gain and its effect with insulin. Insulin was administered intravenously. Fig leaves tea was orally administered to four group of rats. Oral administration of fig leaves tea for 15 days exhibited a significant hypoglycemic effect in hyperglycemic rats at different times of a day. Highest hypoglycemic effect of tea was observed at 3 h which shows that leaves of *Ficus carica* possess significant short term hypoglycemic effect and not so significant long term hypoglycemic effect. When comparison was made between two insulin treatments (10U and 5U insulin) with 1mL *Ficus carica* tea, no significant difference was observed with these two doses, which means that when given in combination of fig leaves tea even 50% decrease of dose of insulin did not produce significant difference in blood glucose level.

**Keywords:** diabetes mellitus, fig leaves, hyperglycemia, herbal medicine

---



## Optimisation of Production Condition of Alkaline Protease Using Indigenous *Bacillus subtilis* from Agricultural Soil

Barkat Ali Solangi<sup>a\*</sup>, Hafiz Rub Nawaz<sup>a</sup>, Uzma Nadeem<sup>a</sup>, Beena Zehra<sup>a</sup> and Agha Asad Noor<sup>b</sup>

<sup>a</sup>PCSIR, Leather Research Centre, D-102, South Avenue, SITE, Karachi, Pakistan

<sup>b</sup>Department of Microbiology, University of Sindh, Jamshoro, Pakistan

(received August 6, 2015; revised May 20, 2016; accepted August 15, 2016)

---

**Abstract.** In this study, proteases have been isolated from agricultural soil samples and then cultured by shake flask method. The growth of the *Bacillus subtilis* has been confirmed by microbiological test on the agar plate and skim milk agar in rough, raised and irregular colonies. The yield of the alkaline protease has been optimised by varying the main factors i.e., nitrogen source (peptone, yeast extract, beef extract, casein, ammonium carbonate and urea), carbon source (sucrose, fructose, mannose, lactose, glucose, maltose and starch), incubation period (12, 24, 36, 48, 72, 84 and 96 h), temperature (35, 40, 45, 50, 55, and 60 °C) and salts (potassium sulphate, magnesium sulphate, calcium sulphate and manganese sulphate). The results revealed that the maximum enzyme production was obtained using casein and minimum activity was obtained using urea as a nitrogen source. Similarly, other factors have shown significant effect on the activity of the enzyme.

**Keywords:** proteases, agricultural soil, shake flask method, optimisation

---

## Deforestation Drivers in Tehsil Barawal, Dir Upper, Pakistan

Anwar Sajjad<sup>a\*</sup>, Syed Adnan<sup>b</sup> and Ahmad Hussain<sup>c</sup>

<sup>a</sup>Department of Environmental Sciences, University of Haripur, KPK, Pakistan

<sup>b</sup>Institute of Geographical Information Systems, National University of Sciences and Technology (NUST), Islamabad, Pakistan

<sup>c</sup>Department of Forestry and Wildlife, University of Haripur, KPK, Pakistan

(received January 14, 2016; revised February 22, 2017; accepted February 23, 2017)

---

**Abstract.** The study area Tehsil Barawal, district Dir (U) of forest has facing serious problems of deforestation. The forest of the study area is ever green forest trees like (*Cedras deodar*, *Abies pindrow*, *Pinus roxbergii*, *Picea smithiana* and *Pinus wallichiana*) are dominate trees. The focus of the study area was to explore the various drivers of the deforestation. The study was based on field survey and GIS and remote sensing techniques. The results of field survey revealed that there are seven drivers of deforestation, source of income from forest, fuel forest collection, illiteracy rate, agriculture expansion, forest fire, illegal cutting/harvesting and encroachment. The field survey shows that 40% people agreed that the major driver of deforestation in the area is agriculture. Study was based on the GIS and remote sensing techniques to explore the various drivers of the deforestation from year 2000 and 2012. In this study the images of 2000 and 2012 were downloaded from landsat 5 satellite. Five key classes such as forest, agriculture, barren land, snow and water were acknowledged. The result showed that the study area Tehsil Barawal, District Dir (U) of forest, barren land, agriculture, water and snow in year 2000 was 49.54, 43.38, 5.19, 1.40 and 0.49%, and the area in 2012 was 37.17, 41.36, 12.69, 5.05 and 3.72%, respectively. After data analysis it was cleared that 2.02% decrease in barren land, 12.37% decrease in forest and 7.5% increase in agriculture land was identified. Therefore, from field as well as from GIS and RS results it was cleared that agriculture is the key driver of the deforestation study area.

**Keywords:** sensing techniques, deforestation, harvesting, Upper Dir.

---

## Population Density Estimates of Punjab Urial in Scrub Forest

Muhammad Irshad Arshad<sup>a\*</sup> and Anwar Hussain<sup>b</sup>

<sup>a</sup>Department of Forestry, Range & Wildlife, Ghazi University, Dera Ghazi Khan, Pakistan

<sup>b</sup>Punjab Wildlife Department, Lahore, Pakistan

(received May 25, 2016; revised May 5, 2017; accepted May 16, 2017)

---

**Abstract.** A ground survey of the population of Punjab urial (*Ovis orientalis punjabiensis*) was conducted in 2007 on transect lines following the distance sampling methodology in five localities of district Jhelum, Punjab, Pakistan. Its aim was to determine the population status with reference to trophy sized males. The estimated population density for 5 distinct survey areas (Kandal Un-classed Forest and Jalapur Wildlife Sanctuary, Phadial Reserved Forest, Nili South and Brali Reserved Forests, Lehri and Jindi Reserved Forests and Tilla Jogian Game Reserve). Per square kilometer of area, respectively was  $1.26 \pm 0.49$ ,  $1.30 \pm 0.56$ ,  $0.46 \pm 0.19$ ,  $0.36 \pm 0.15$  and  $0.37 \pm 0.13$ , and overall for combined data it was  $0.63 \pm 0.14$ . The male/female ratio indicated a preponderance of females over the males (1:2.53). The estimated mean group size was  $5.08 \pm 0.44$ . The study showed some indications of recovery in the population of Punjab urial in a context of increased protection efforts by the Punjab Wildlife Department in the recent past by the introduction of community-based trophy hunting in the area as a conservation and management tool.

**Keywords:** Punjab urial, *Ovis vignei punjabiensis*, distance sampling, line transect, population density

---

# Comparative Toxicity of Spinetoram and Nitenpyram against Earthworm and their Effects on Protein Contents and Cholinesterase Activity

Altaf Hussain<sup>a\*</sup>, Muhammad Farhanullah Khan<sup>a</sup>, Omer Mukhtar<sup>b</sup>, Muhammad Faheem<sup>a</sup>  
and Umeed Ali<sup>b</sup>

<sup>a</sup>Department of Zoology, University of Karachi, Karachi-75270, Pakistan

<sup>b</sup>Food & Marine Resources Research Centre, PCSIR Laboratories Complex, Karachi-75280, Pakistan

(received December 19, 2016; revised July 29, 2017; accepted August 1, 2018)

---

**Abstract.** The earthworms' numbers and biodiversities are considered as an indicator of soil nature and fertility. Earthworms are very important soil organisms that aid in the decomposition of plant litter. In recent era earthworm obtain more attention in agriculture practices, so that in the present work to determined the comparative effects of nitenpyram and spinetoram on total amount of protein and cholinesterase activity of *Pheretima posthuma* Kinberg (Order: Haplotaxida and Family: Megaseolecida ) earthworm. Total protein contents were found to be decreased significantly in the peristomium, clitellum, and abdomen regions under treatments of nitenpyram and spinetoram. Cholinesterase activities were inhibited in the peristomium, clitellum and abdomen under the exposures of nitenpyram and spinetoram, at the intervals of 30, 60 and 90 seconds in *P. posthuma*. Conclusively it is safe to say that spinetoram is better than neonicotinoids for the earthworms, therefore could be switched from neonicotinoids as an IPM component.

**Keywords:** enzyme, bio-pesticide, neonicotinoid, earthworm

---

## Short Communication

# Correlation and Regression Coefficient Estimates between some Growth Performance Traits of Harnai Sheep

Rameez Raja Kaleri<sup>a\*</sup>, Hubdar Ali Kaleri<sup>a</sup>, Asma Kaleri<sup>a</sup>, Abdul Rashid<sup>b</sup>, Rashid Ali Shah<sup>a</sup>, Gul Muhammad Marri<sup>c</sup>, Rajesh Kumar and Deepesh Kumar<sup>a</sup>

<sup>a</sup>Department of Animal Breeding & Genetics, Sindh Agriculture University, Tandojam, Sindh, Pakistan

<sup>b</sup>Mountain Agriculture Research Institute (PARC), Skardu, Gilgit, Pakistan

<sup>c</sup>Arid Zone Research Institute (PARC), Umarmkot, Sindh, Pakistan

(received April 13, 2017; revised June 16, 2017; accepted July 18, 2018)

---

**Abstract.** Present investigation was carried out to estimate the correlation and regression coefficient estimates between some growth performance traits of Harnai sheep. The available data was collected and maintained at Multipurpose Sheep Research Centre Yetabad, District Loralia, Balochistan, Pakistan. The data regarding growth traits including birth weight, yearling weight, weaning weight and fleeces weight was analysed for the estimation of correlation and regression coefficient. In present study the results for correlation and regression coefficient estimates were observed low to positive between some growth performance traits of Harnai sheep. Due to low results for correlation and regression coefficient, it is concluded that mass selection is advisable for achieving better performance in growth traits of Harnai sheep.

**Keywords:** correlation, regression coefficient, growth traits, Harnai sheep

---

## Review

# A Review on Rangeland Management in Pakistan, Bottlenecks and Recommendations

**Muhammad Jamil<sup>a</sup>, Muhammad Mansoor<sup>a\*</sup>, Fawad Anwar<sup>a</sup>,  
Sher Muhammad<sup>b</sup> and Aftab Ahmed Awan<sup>a</sup>**

<sup>a</sup>Arid Zone Research Centre (AZRC, PARC), Dera Ismail Khan-29050, Pakistan

<sup>b</sup>National Agricultural Research Center, Islamabad, Pakistan

(received August 25, 2016; revised March 30, 2017; accepted May 2, 2017)

---

**Abstract.** Role of rangelands in the livelihoods improvement of rural communes' is pivotal, through animal rearing, obtaining versatile products and services. Rangelands play key role in maintenance of agro-ecosystem and preservation of biodiversity in the country by improving infiltration rate, leading thereby a sustainable water flow in the down streams and reducing soil erosion. However, due to increased population pressure there is analogous increase in the demand for food, forage and other resources. In addition rangelands provide wildlife and fish habitat and recreation spots. Keeping in view the current global climate change scenario the government has given high priority to rangelands management, hence the concentration of the politicians and policy makers is to focus on the sustainable rangeland development and management. The article deals first with the importance of range management, then discusses rangelands problems of Pakistan, causes of degradation of rangelands, human factors contributing to range degradation and finally recommendations are discussed.

**Keywords:** Pakistan, rangeland, livestock, management, biodiversity

---

# INSTRUCTIONS TO AUTHORS

*Pakistan Journal of Scientific and Industrial Research, Series A: Physical Sciences* and *Series B: Biological Sciences* publish original researches as **full research papers, short communications and reviews**. Articles submitted to other journals for publication are not accepted. All manuscripts are evaluated by subject experts of advanced countries. However, authenticity of the contents remains the sole responsibility of the author/s. The Journal reserves the rights to edit, wherever considered necessary.

Manuscripts should be submitted with a covering letter, giving unabbreviated names and addresses of all the authors, indicating the corresponding author and certifying that the paper has neither been, nor will be sent elsewhere for publication.

Author/s may submit electronic copies in MS Word via e-mail or through our website. The manuscript should contain, in order, Running Title (not more than 60 characters), Title of the article, full names, addresses, telephone numbers, fax numbers, e-mail addresses of all the Authors, Abstract (not more than 150 words), three to six Keywords, Introduction, Materials and Methods, Results and Discussion, Acknowledgement (if applicable), References, Tables and Illustrations. All illustrations should be of good quality with legible captions. In case of coloured figures, 50% printing expenses would have to be borne by the authors. IUPAC rules for units and their abbreviations should be used. **Plagiarism Report** of the submitted paper must be attached and similarity index should be  $< = 19\%$ .

References should be verifiable and cited in the text by the last name of the author (both authors if only two, the first author and *et al* when more than two) followed by the year, **in descending chronological order**. All references in the bibliography should be listed in alphabetical order of the authors' last names followed by date of publication and other complete details with full names of Journals as given below. Anonymous (Anon) references are not accepted.

## Papers of Journals

### In "Text"

Park (2005), Aksu and Kabasakal (2004) and Evans *et al.* (2000) (Park, 2005; Aksu and Kabasakal, 2004; Evans *et al.*, 2000)

### In "References"

Evans, W.J., Johnson, M.A., Fujimoto, Cy. H., Greaves, J. 2000. Utility of electrospray mass spectrometry for the characterization of air-sensitive organolanthanides and related species. *Organometallics*, **19**: 4258-4265.

### Books

Cinar, A., Parulekar, S.J., Undey, C., Birol, G. 2003. *Batch Fermentation: Modeling, Monitoring, and Control*, 250 pp. Marcel Dekker Inc., New York, USA.

### Chapters in Edited Books

Newby, P.J., Johnson, B. 2003. Overview of alternative rapid microbiological techniques. In: *Rapid Microbiological Methods in the Pharmaceutical Industry*, M.C. Easter (ed.), vol. **1**, pp. 41-59, 1<sup>st</sup> edition, Interpharm/CRC, Boca Raton, Florida, USA.

### Papers in Proceedings

Marceau, J. 2000. Innovation systems in building and construction and the housing industry in Australia. In: *Proceedings of Asia-Pacific Science and Technology Management Seminar on National Innovation Systems* pp. 129-156, Japan Int. Sci. Technol. Exchange Centre, Saitama, Japan.

### Reports

*SIC-PC SIR. 2002. Biannual Report, 2000-2001; 2001-2002*, Scientific Information Centre, Pakistan Council of Scientific and Industrial Research, PCSIR Laboratories Campus, Shahrah-e-Dr. Salimuzzaman Siddiqui, Karachi, Pakistan.

### Thesis

Saeed, A. 2005. Comparative Studies on the Biosorption of Heavy Metals by Immobilized Microalgal Cultures, Suspended Biomass and Agrowastes. *Ph.D. Thesis*, 248 pp., University of the Punjab, Lahore, Pakistan.

### Patents

Young, D.M. 2000. Thermostable Proteolytic Enzymes and Uses Thereof in Peptide and Protein Synthesis, US Patent No. 6,143,517, 7<sup>th</sup> November, 2000.

**Data Set from a Database** (information should be retrievable through the input)

Deming, D.; Dynarski, S. 2008. The lengthening of childhood (NBER Paper 14124) Cambridge, MA: National Bureau of Economic Research. Retrieved July 21, 2008 from <http://www.nber.org/papers/w14124> DOI No.....

**The manuscripts should be addressed to: The Executive Editor, Pakistan Journal of Scientific and Industrial Research, Scientific Information Centre, PCSIR Laboratories Campus, Sharah-e-Dr. Salimuzzaman Siddiqui, Karachi-75280, Pakistan.**

**Phone:** (92-21)-99261914, 16, 17, 99261949,

**Fax:** (92-21) 99261913

**Web:** <http://www.pjsir.org>; **E-mail:** [info@pjsir.org](mailto:info@pjsir.org)