

Short Communications

Pakistan J. Sci. Ind Res., Vol. 19, Nos. 3-4, June-August 1976

UTILIZATION OF N BY WHEAT FROM NH₄ AND NO₃ FORMS APPLIED AS NH₄NO₃ SELECTIVELY LABELLED WITH N¹⁵

A. HAMID

Atomic Energy Agricultural Research Centre, Tandojam

G. SARWAR

Nuclear Institute for Agriculture and Biology, Lyallpur

(Received May 5, 1975; revised December 11, 1975)

Uptake of NH₄ and NO₃ ions by plants has been studied extensively.¹⁻³ When nitrification inhibitor was used to differentiate between the ions being absorbed by wheat NH₄-N promoted more leaf and stem growth than NO₃-N during early stages but the grain yield was unaffected by the form of N.³⁻⁵ The economics of fertilizer use requires an assessment of the utilization of N from various sources. Isotopically labelled fertilizer materials provide a unique tool for such investigations. Fried and Dear⁶ suggested an A value concept for measuring the amount of nutrient in a source in terms of a standard source of that nutrient. It is based on the assumption that when a plant has access to two sources of N with no interaction between the sources, the plant will absorb N from the two sources in direct proportion to the respective amounts available.

The objective of this study was to determine the utilization of N from NH₄ and NO₃ in wheat from NH₄NO₃ selectively labelled with N¹⁵ and applied at seeding and at tillering stage.

Materials and Methods

A field experiment was done on sandy clay-loam soil having pH 8.5, total N 0.03% and available P

(Olson) 10 p.p.m. Fertilizer treatments were arranged in a randomized block design with eight replicates. The individual plot was 1 × 5 m with five rows of wheat plants. Ammonium nitrate was given in two doses of 60 kg N/ha each applied at seeding and tillering stage. Each plot received only one dose of N from selectively labelled NH₄NO₃ (N¹⁵H₄NO₃ or NH₄N¹⁵O₃ with 1% N¹⁵ enrichment) either at seeding or at tillering. Superphosphate was applied at 30 kg P/ha at seeding. At seeding both NH₄NO₃ and superphosphate fertilizers were applied in a band 5 cm to the side and 2 cm below the seed row. At tillering stage NH₄NO₃ was broadcast uniformly on the soil. The crop received normal irrigation.

Grain and straw samples were collected at maturity, dried at 70°C, ground and analysed for N content by Kjeldahl method. The N¹⁵ analysis was done by the staff at Sheibersdorf Laboratories of IAEA.

The efficiency of uptake of NH₄-N and NO₃-N was evaluated by A value data calculated from N¹⁵ analysis of the grain and based on the formula.⁶

$$A=B (1-Y/Y)$$

where A (A value) is the amount of soil N measured in units of applied fertilizer standard; B is the amount of labelled standard N fertilizer applied; and Y is the fraction of the N in the plant derived from the labelled standard N fertilizer (1-Y, thereby the fraction from the soil).

An analysis of variance was made and L.S.D. worked out to test the significance of differences between means.

Results and Discussion

The results showed that the time of application of N had a significant effect on uptake of fertilizer N in wheat grain. The utilization of both NH₄-N and NO₃-N from NH₄NO₃ applied at tillering stage was greater than that from the dose applied at seeding (Table 1). This indicates that tillering is the critical stage of wheat for the application of N and at this stage N is most efficiently utilized under conditions prevailing in Pakistan. Lower recovery of N applied

TABLE 1. EFFECT OF NH₄NO₃ ON YIELD, N UPTAKE, % N DERIVED FROM FERTILIZER, UTILIZATION OF FERTILIZER N AND 'A' VALUE IN WHEAT.

Stage of N application kg N/ha		Yield kg/ha		Total N uptake kg/ha		% N derived from fertilizer		Fertilizer N uptake kg/ha		Utilization of fert. N %		A value kg/ha		
(Seeding)	(Tillering)	(Grain)	(Straw)	(Grain)	(Straw)	(Grain)	(Straw)	(Grain)	(Straw)	(Grain)	(Straw)	(Grain)		
N¹⁵H₄NO₃														
60*	60	4704	8264	88.9	27.6	9.4	8.6	8.4	2.8	28.0	8.0	289		
60	60*	4598	8074	87.4	25.3	14.0	14.1	12.4	3.9	41.3	13.0	184		
NH₄N¹⁵O₃														
60*	60	4604	8097	87.0	24.7	11.6	13.4	10.3	3.3	34.0	11.0	226		
60	60*	4699	8108	88.2	26.1	17.9	18.4	15.7	4.5	52.3	15.0	137		
L.S.D.		P=0.05 P=0.01		NS	NS	NS	NS	6.2 8.2	7.0 9.1	2.44 3.55	NS	6.00 11.06	NS	47 70

*Fertilizer labelled with N¹⁵. NS nonsignificant.

at seeding may be because a part of N was leached down and distributed in deeper layers and wheat seedlings with less developed root system were unable to absorb N from soil layers beyond the root system. The comparison of utilization of N in wheat grain from NH_4 and NO_3 forms showed that wheat plants absorbed 21.4 and 26.6% higher N from NO_3 than from NH_4 form from NH_4NO_3 carrier applied at seeding and tillering stage, respectively ($P=0.05$). Similarly per cent utilization of N in straw from NO_3 form was greater than from NH_4 form. Other criterion used for assessing the efficiency of fertilizer N like A value also indicated that NO_3 form of N was significantly superior to NH_4 form for wheat. Lower A value is an indicator of higher utilization of applied fertilizer by the plants. The smaller recovery of N from NH_4 than from NO_3 may be because a sizeable part of NH_4 was lost through volatilization under alkaline soil conditions and chemical immobilization of NH_4 in the soil. 8, 9

The results of this study are in agreement with those of Spratt and Gasses¹⁰ who reported that wheat plants given NO_3 contained significantly more N than those given NH_4 when watered adequately. The results obtained in this study and those reported earlier⁷ suggest that NO_3 source of N would be a better carrier for irrigated wheat under Pakistan conditions.

Acknowledgement. This work was done under IAEA contract No. 639/R2/RB. The N^{15} -labelled fertilizers were proved by joint FAO/IAEA Division of Atomic Energy in Food and Agriculture. Grateful acknowledgement is extended to IAEA for analysis of plant samples for N^{15} . Thanks are due to Mr. Farooq Maqsood, Statistician, Nuclear Institute for Agriculture and Biology, Lyallpur, for statistical analysis of data.

References

1. E. J. Hewitt in *Nitrogen Nutrition of the Plant*, edited by E. A. Kirkby (University of Leeds Press, Leeds, 1970), p. 220.
2. F. G. Viets Jr. in *Soil Nitrogen. Agronomy* (edited by W. V. Bartholomew and F. E. Clark) (Amer. Sec. Agron., Madison, Wis., 1955), 10, 10, p. 503.
3. E. D. Spratt and J. K. R. Gasser, *J. Agr. Sci.*, **74**, 111 (1970).
4. E. D. Spratt, *Soil Sci. Soc. Am. Proc.*, **37**, 259 (1973).
5. E. D. Spratt, *Agron. J.*, **66**, 57 (1974).
6. M. Fried and L. A. Dean, *Soil Sci.*, **73**, 263 (1952).
7. A. Hamid, *Plant Soil*, **37**, 389 (1972).
8. S. G. Misra and B. Singh, *Indian J. Agron.*, **14**, 214 (1969).
9. E. D. Spratt and J. K. R. Gasser, *Can. J. Soil Sci.*, **50**, 263 (1970).
10. E. D. Spratt and J. K. R. Gasser, *Can. J. Pl. Sci.*, **50**, 613 (1970).

Pakistan J. Sci. Ind. Res., Vol. 19, Nos. 3-4, June-August 1976

APPEARANCE OF A POTENTIAL PEST OF RICE, *PODOPS LIMOSA* WALKER (PENTATOMIDAE: PODOPINI) THE STINK BUG OF PADDY, IN THE RICE FIELDS OF SUJAWAL, THATTA DISTRICT OF LOWER SIND, PAKISTAN*

IMTIAZ AHMAD and MOHAMMAD AFZAL

Department of Zoology, University of Karachi

(Received November 3, 1975)

Podops limosa Walk. the stink bug of paddy, is closely allied to *Scotinophara coarctata* (Fabr.) which has been reported as major pest of rice in the Salem district of former British India,² Malaysia,¹ Sarwak⁴ and Thailand.

During a recent survey of the insect pests of paddy throughout southern Sind, *P. limosa* has only been found in paddy fields of Sujawal in annoying numbers. (Actual population studies are still being carried out and will be reported on separately.)

The stink bug invades only those fields which have no standing water and sucks the sap from paddy stem, as a result of which panicles fail to develop, the leaves turn reddish brown and sometimes the whole plant is stunted. All developmental stages of the bug devour on the paddy plant.

If an infested field is flooded both nymphs and adults escape to adjacent sedges and grasses, some of which probably serve as alternate host.

At this stage it has been observed that drying up of a paddy field may invite these bugs in large numbers even a few days before harvest.

As regards the taxonomic position of the bugs it could be tentatively suggested that a complex of species are involved. Detailed systematic and biological studies are already in progress which will further throw light on the present complex.

References

1. G. H. Corbett and M. Yusupe, *Malayan Agr. J.*, **12**, 91. (1924).
2. H. M. Lefroy, *Indian Insect Life*. London. 1909.
3. T. Wongsiri, K. Kovitvadhi, *proc. Symp. Int;ng Rice Res. Inst.*, 571 (1964).
4. A. Yunus and H. G. L. Rowchild, *Proc. Symp. Intern. Rice Inst.*, 635 (1964).

*Financially supported by a PL 480 Res. Project. No. FG. Pa 181 (A 17. ENT 37).

Pakistan J. Sci. Ind. Res., Vol. 19, Nos. 3-4, June-Aug. 1976

A NEW SPECIES OF AMBLYTHYREUS WESTWOOD (HETEROPTERA, REDUVOI DEA, PHYMATIDAE) FROM PAKISTAN

NARJIS YOUSUF and IMTIAZ AHMAD

Department of Zoology, University of Karachi

(Received November 3, 1975)

Oriental phymatid fauna is rich yet unexplored, although it is an important group of predatory insects.

Uhler² found some species on the stems and flowers of Euphorbiaceae and Compositae praying upon visiting bees and other insects.

Distant¹ described six species of *Amblythyreus* from Assam, Burma and Bengal, mostly on colouration, ratio of the length of antennal segments, length and width of pronotum and scutellum.


The described unique female specimen has bizarre rows of spines on the median portion of the pronotum, three of these located in front of depressed line demarcating anterior half from posterior half and remarkably raised outer margins of the latter as well as several prominent characters noted in the description. The above characters distinguish this specimen as a new species *A. murreeana*.

Amblythyreus murreeana n.sp. (Figs. 1-3)

Body ovate, tylus, jugae, vertex, antennal segments, pronotum (except ridge), scutellum, abdominal sternites, labial segments and legs brown. Connexiva, wing membrane and eyes ochraceous, ocelli bright red.

Head short, but distinctly longer than broad, with anterior notch, densely granulated, posterior portion (including eyes) more than three times as wide as anterior portion of head (excluding eyes), with three pairs of lateral spines. Jugae not meeting, pointed at the apices with a pair of spines, anteriorly produced with deep median notch, tylus much smaller than jugae, eyes very slightly convex, inserted in anterior half of the head. Ocelli on mounts facing each other, located in the centre, lateral margins of posterior portion of head with three pairs of spines evenly spaced, antenniferous tubercles withholding lateral half of basal antennal segments, latter incrassated and granulated, apical segments leaf-shaped (Fig. 1a), labium reaching anterior coxae with II, III and IV segments almost equal in length, length of head, 0.433 mm; length of anterior portion (excluding eyes), 0.1 mm.; of posterior portion (including eyes) 0.333 mm; width, 0.333 mm; interocular distance, 0.3 mm; interocellar distance, 0.2 mm; length of antennal segments I, 0.1 mm; II, 0.64 mm; III, 0.066 mm. Pronotum distinctly broader than long, strongly narrowing, anterior margin four times narrower than width across posterior margin, former shorter than width of head across eyes, deeply concave forming socket for posterior portion of head, surface of anterior half densely granulated, lateral margins slightly dentate, posterior half densely punctate, dumb-bell-shaped, lateral margins sinuate, humeral angles acute, distinctly elevated, posterior margin sinuate, five pairs of spines arising from a central ridge, located anterior to a depressed line, demarcating anterior and posterior half, fifth pair larger and located more anteriorly, spines subequal in length, tooth-shaped, width of pronotum, 1.5 mm.; length 1.0 mm.

Scutellum short, heart-shaped, attaining fourth abdominal segment, punctate, pointed apically, slightly raised area at the base continued medially, bordered with yellow granules, length, 0.7 mm.; width, 0.7 mm.;



Figs. 1-3

distance apex scutellum apex abdomen, 1.5 mm. Metathoracic scent gland ostioles, slit-like, peritremes fully developed, spout-like with spines broadly rounded and projecting postero-laterad, reaching outer limit of the evaporatoria, latter prominent and rugulose.

Membrane of hemelytra extended beyond apex of abdomen, with simple longitudinal veins, clavus and corium exposed, abdomen much expanded laterally and flattened, longer than broad, connexiva completely exposed, sternites medially projected anteriorly fitting into concavity on posterior margin of segment III and VI.

Forelegs modified for predation with femur flattened, broad, distal margin grooved, to receive small curved toothed tibia (Fig. 2).

Posterior margin of seventh abdominal sternum concave, eighth paratergites greatly developed, lobe-like with acute apices, base broad, extending much beyond rather smaller ninth paratergites, first gonocoxae much reduced, telescoped within eighth paratergites, proctiger U-shaped, enlarged (Fig. 3).

Comparison. *Amblythyreus murreeana* new species is closely related to *A. gestroi* but can easily be separated from the latter by the acute and elevated humeral angles of pronotum with five pairs of spines arising from a ridge in the middle of pronotum, three of these located in front of a depressed line demarcating the anterior from the posterior half. In the new

species, on the base of scutellum there is a slightly elevated area which is continued medially but is absent in *A. gestroi*. The latter has a small spot at each basal angle of the scutellum, being absent in *A. murreana* new species.

Material Examined. Holotype, female, Pakistan: Punjab, Murree Hills (Sandian village), on *Rumex hastatus* leg. Azhar A. Khan, 25 May 1971, lodged at Natural History Museum, Department of Zoology,

University of Karachi.

References

1. W. L. Distant. *The Fauna of British India including Ceylon and Burma*, Rhynchota, vol. II, London, 1904).
2. P. R. Uhler, Bull. U.S. Geol. Geog. Surv. Terr., 3, 355, 765 (1877).