

## CURRENT STATUS OF NICKEL CONTAMINATION OF LOCAL VEGETABLE GHEE

Sadaf Yaqub, M Jaffar\* and Khalid Masud

Department of Chemistry, Quaid-i-Azam University, Islamabad, Pakistan

(Received 11 July 1988; accepted 1 December 1998)

Nickel contamination of locally produced vegetable ghee has been evaluated using the atomic absorption flame photometric method, employing n-hexane as solubilization medium. The data, based on 3 batches of samples comprising a total of 32 samples procured from 18 manufactures, is reported as  $\bar{X} \pm SD$ , for 3 to 5 replicates. The nickel concentration ranges for the three batches are, 0.445-0.726 mg 100 g<sup>-1</sup>, 0.668-1.2668 mg 100 g<sup>-1</sup>, and 0.395-1.234 mg 100 g<sup>-1</sup>, thereby indicating an inefficient nickel filtration process at individual industrial units. Substantial comparative variations in the nickel concentration are observed in the output of a given ghee industry. The estimated levels of nickel exceed the WHO's lower safe limit of 0.10 mg 100 g<sup>-1</sup> ghee serving.

**Key words:** Nickel contamination, Vegetable ghee, Hydrogenation.

### Introduction

Nickel is well known for its toxic effects on human health, which range from minor to severe pathological pulmonary disorders (Schroeder *et al* 1962 and Nielsen 1993). It is also well documented that nickel toxicity is associated with lipid peroxidation in the target organ (Dabeka and McKenzie 1995 and WHO 1989). The health hazards of nickel have thus caused concern among health authorities all over the world (WHO 1991a and WHO 1991b). In view of this, the daily total intake of nickel from various foods is laid down between 100-400 µg (Macdonald 1991 and Parr 1992). In developed countries, stringent control over the level of the metal during ghee manufacture is being legally regulated and exercised. Unfortunately, in developing countries, the state of affairs is far from being satisfactory. Nickel, used as a catalyst in the hydrogenation of edible oils for the manufacture of ghee, is subsequently removed from the reaction medium by filtration. The metal, however, permeates at various trace levels into the final product, ghee, which is a such abundantly used as a cooking medium by a large segment of Pakistan population (Dean and Ayub *et al* 1993).

The present study was undertaken to assess the present status of nickel contamination of locally produced vegetable ghee so as to help law enforcing and health authorities to evolve a control policy for decontamination of this toxic metal from an essential edible commodity.

### Material and Methods

Three batches of ghee samples were procured from local retailers, comprising of 19, 8 and 5 samples respectively

(Table 1). All samples were collected in new, strong polythene bags of 1 kg capacity. Similarly, 7 edible oil samples (Table 3) were procured for the estimation of background nickel content. The solubilization medium for the samples was n-hexane which is well known for its efficient and stable thermal combustion characteristics and capability for improved flow rate of sample solution during aspiration. Quantification of results was based on standard calibration method, using a Shimadzu atomic absorption spectrophotometer (model AA-670), duly backed by automated background compensation facility. An exactly weighed (preferably 5.000 g) ghee/oil sample was transferred to a pre-washed and pre-dried 30 mL capacity centrifuge tube containing 20.0 mL of carefully measured n-hexane. The tube was given a swirling motion for about 1 minute, placed in the centrifuge rack and spun around at 3000 rpm for five minutes. After separating the settleable insoluble matter, the clear solution containing dissolved ghee/oil, which contained nickel in a dispersed form, was aspirated directly with the following operational condition:  $\lambda = 232.0$  nm; HCL current = mA; slit width = 0.15 nm; fuel (acetylene) flow rate = 0.9 L<sup>-1</sup>min. The standards, within a range of 1-5 µg g<sup>-1</sup> nickel, were prepared from nickel phthalocyanine dye (FW 1148.11, 99% pure Merck) dissolved in hexane. The detection limit achieved was 0.010 ppm at x 10 expansion. Three to five replicates were simultaneously run and the results are reported as  $\bar{X} \pm SD$ .

### Results and Discussion

The results of various batches of ghee samples (Table 2) showed a substantial variability of nickel concentration in individual samples. The metal concentration was found to range from 0.445 mg 100g<sup>-1</sup> to 0.726 mg 100g<sup>-1</sup> for batch 1; from 0.668 mg

\*Authors for correspondence


**Table 1**  
Sample codes and description of various vegetable ghee brands/batches analyzed for nickel

Serial No.	Sample Code/Batch	Brand	Manufacturer
1.	S-1/1, S-1/2 S-1/3	Mujahid Banaspati Fauji Banaspati	Hafeez Iqbal Oil & Ghee Industries Waheed Hafeez Ghee Industries (Pvt) Ltd.
2.	S-2/1, S-2/2 S-2/3	Family Banaspati	Pan Asia Food Products (Pvt) Ltd. Tahir Oil Mills
3.	S-3/1, S-3/2, S-3/3	Dalda Banaspati	Lever Brothers Pakistan Ltd.
4.	S-4/1, S-4/2 S-4/3	Fauji Banaspati Sawaira Banaspati	Fauji Industries (Pvt) Ltd. Sawaira Ghee Industries
5.	S-5/1, S-5/2 S-5/3	Tullo Banaspati ACP Banaspati	Wazir Ali Industries Ltd. ACP Oil Mills (Pvt) Ltd.
6.	S-6/1, S-6/2	Zaiqa Banaspati	Punjab Oil Mills (Pvt) Ltd.
7.	S-7/1 S-7/2	Kohinoor Banaspati Asia Banaspati	Silver Oil Mills (Pvt) Ltd. Pan Asia Foods (Pvt) Ltd.
8.	S-8/1 S-8/2	ACP Banaspati Fauji Banaspati	Waheed Hafeez Ghee Industries (Pvt) Ltd.
9.	S-9/1	Anarkali Banaspati	Fazal Vegetable Ghee Mills
10.	S-10/1	Latif Banaspati	ACP Oil Mills (Pvt) Ltd.
11.	S-11/1	Asia Banaspati	Pan Asia Food (Pvt) Ltd.
12.	S-12/1	Haseeb Banaspati	ACP Oil Mills (Pvt) Ltd.
13.	S-13/1	Habib Banaspati	ACP Oil Mills (Pvt) Ltd.
14.	S-14/1	Kashmir Banaspati	United Industries Ltd.
15.	S-15/1	Kissan Banasopati	Kissan Ghee Corporation (Pvt) Ltd.
16.	S-16/1	Kauser Banaspati	Kauser Ghee Mills (Pvt) Ltd.
17.	S-17/1	POF Japan Banaspati	POF Ghee Industries
18.	S-18/1	GCP Banaspati	Ghee Corporation Pakistan Ltd.
19.	S-19/1	Kiran Banaspati	Kiran Ghee Industries

100g<sup>-1</sup> to 1.268 mg 100g<sup>-1</sup> for batch 2 and from 0.395 mg 100g<sup>-1</sup> to 1.234 mg 100g<sup>-1</sup> for batch 3. Maximum concentration was 1.268 mg 100g<sup>-1</sup> (batch 2), 0.726 mg 100g<sup>-1</sup> (batch 1) and 1.234 mg 100g<sup>-1</sup> (batch 3) in ghee manufactured by Asia Foods, Tullo Wazir Ali Industries and Dalda Lever Brothers, respectively. The results indicated that even the leading ghee manufacturers were unable to cope with stringent quality control regarding nickel content of their finished product. On the whole, the average nickel content for batch 1 ghee samples was 0.589 mg 100g<sup>-1</sup> for batch 2, the average concentration was 0.872 mg 100g<sup>-1</sup> and for batch 3, it was 0.699 mg 100g<sup>-1</sup>. Batch 3 was found to contain maximum nickel; both reputed and non-reputed ghee manufacturers share this reputation.

The World Health Organization (WHO Report 1993) has laid down a lower safe limit of acceptable nickel content as 0.1 mg 100g<sup>-1</sup> serving of vegetable ghee which, in the present case, enhanced to the level of 0.5 mg 100g<sup>-1</sup>. Only 7 ghee samples qualified for being acceptable from the viewpoint of safe ingestion (Table 2); thus, about 78% of the ghee brands contained unacceptable nickel level.

The results indicate that the current situation of nickel content of ghee is not satisfactory for consumers. A comparative evaluation of this situation is illustrated in Fig. 1. For example, in Dalda Banaspati the nickel content of batch 1 was almost 50% lower than that in batch 2, and it further increased in batch 3. Similar variations are also observed in other brands of ghee.



Comparative levels of nickel in various batches of ghee samples from a given manufacturer. (-----, Limit recommended by WHO.)

**Table 2**  
Levels of nickel ( $\bar{X} \pm SD$ , mg 100g<sup>-1</sup>) in various vegetable ghee samples

S.No.	Sample Code/ Batch 1	Ni Level	S.No.	Sample Code/ Batch 2	Ni Level	S.No.	Sample Code/ Batch 3	Ni Level
1.	S-1	$\bar{X}$ 0.667 $\pm SD$ 0.012	20	S-1	$\bar{X}$ 0.687 $\pm SD$ 0.013	28	S-1	$\bar{X}$ 0.514 $\pm SD$ 0.010
2.	S-2	$\bar{X}$ 0.640 $\pm SD$ 0.014	21	S-2	$\bar{X}$ 0.783 $\pm SD$ 0.009	29	S-2	$\bar{X}$ 0.339 $\pm SD$ 0.007
3.	S-3	$\bar{X}$ 0.445 $\pm SD$ 0.016	22	S-3	$\bar{X}$ 1.083 $\pm SD$ 0.014	30	S-3	$\bar{X}$ 1.234 $\pm SD$ 0.011
4.	S-4	$\bar{X}$ 0.535 $\pm SD$ 0.009	23	S-4	$\bar{X}$ 1.083 $\pm SD$ 0.016	31	S-4	$\bar{X}$ 0.657 $\pm SD$ 0.009
5.	S-5	$\bar{X}$ 0.597 $\pm SD$ 0.013	24	S-5	$\bar{X}$ 1.268 $\pm SD$ 0.016	32	S-5	$\bar{X}$ 0.697 $\pm SD$ 0.010
6.	S-6	$\bar{X}$ 0.556 $\pm SD$ 0.012	25	S-6	$\bar{X}$ 0.710 $\pm SD$ 0.011			
7.	S-7	$\bar{X}$ 0.539 $\pm SD$ 0.009	26	S-7	$\bar{X}$ 0.690 $\pm SD$ 0.012			
8.	S-8	$\bar{X}$ 0.502 $\pm SD$ 0.007	27	S-8	$\bar{X}$ 0.068 $\pm SD$ 0.013			
9.	S-9	$\bar{X}$ 0.571 $\pm SD$ 0.008						
10.	S-10	$\bar{X}$ 0.495 $\pm SD$ 0.010						
11.	S-11	$\bar{X}$ 0.726 $\pm SD$ 0.010						
12.	S-12	$\bar{X}$ 0.531 $\pm SD$ 0.005						
13.	S-13	$\bar{X}$ 0.609 $\pm SD$ 0.008						
14.	S-14	$\bar{X}$ 0.495 $\pm SD$ 0.011						
15.	S-15	$\bar{X}$ 0.494 $\pm SD$ 0.013						
16.	S-16	$\bar{X}$ 0.609 $\pm SD$ 0.012						
17.	S-17	$\bar{X}$ 0.501 $\pm SD$ 0.012						
18.	S-18	$\bar{X}$ 0.458 $\pm SD$ 0.009						
19.	S-19	$\bar{X}$ 0.569 $\pm SD$ 0.010						

**Table 3**

Code, description and nickel content of various oils collected from local manufactures

S.No.	Sample code	Nickel concentration (mg kg <sup>-1</sup> )	Brand
1.	SO-1	0.061	Rafhan corn oil
2.	SO-2	0.087	Canola cooking oil
3.	SO-3	0.080	Sunflower cooking oil
4.	SO-4	0.069	Dalda cooking oil
5.	SO-5	0.083	Sharlo oil, UAE
6.	SO-6	0.071	Fauji oil (cotton seed)
7.	SO-7	0.090	Habib oil

A total of 7 oil samples, used in the manufacture of ghee, were analyzed to investigate the nickel content originally present in these oils prior to hydrogenation. Irrespective of origin and type of oil, the range of nickel concentration found was between 0.061 mg<sup>-1</sup>kg and 0.090 mg kg<sup>-1</sup> (Roggi *et al* 1997). This, being the background concentration of nickel, could not contribute so heavily to the nickel content of ghee evaluated in the present study.

The majority of the ghee samples were found to contain undesirably high nickel levels as per WHO safe standards. It is high time that a control over the quality of ghee from the viewpoint of its nickel content be promulgated by the local law enforcing authorities.

#### Acknowledgement

We are grateful to the Nutrition Division, National Institute of Health, for providing the facility of inter-laboratory comparison of data using standard reference material.

#### References

- Dabeka R W, McKenzie A D 1995 Survey of lead, cadmium, fluoride, nickel and cobalt in food composites and estimation of dietary intakes of these elements by Canadians in 1996-1988. *J AOAC International* **78** 897-909.
- Dean T H, Ayub M A 1993 *Vitamin A content in banaspati ghee and oil produced in Pakistan*. UNISEF Report Islamabad pp8-15.
- Macdonald I 1991 *Monitoring dietary intakes*. Springer-Verlag, Berlin. pp 90-98.
- Nielsen F H 1993 Ultratrace elements of possible importance for human health: an update. *Prog Clin Biol Res* **380** 355-376.
- Parr R M 1992 *Dietary intakes of trace element*. IAEA Report Vienna Austria.
- Roggi C, Merol E, Minoia C, 1997 Nickel food exposure. *Ann Ig* **9** 221-227.
- Schroeder H A, Balassa J J, Tipton I H 1962 Abnormal trace metals in Man. *Nickel J Chronic Disease* **15** 51-65.
- WHO (Report of expert committee) 1989 *Trace elements in human nutrition*. Geneva Switzerland, Series no.853. pp 143-150.
- WHO Report 1991a *IPCS Nickel*. **108** pp 143-150.
- WHO Report 1991b *Nickel, health and safety guide*. **62** pp 12-20.
- WHO Report 1993 *Recommendations and health criteria for metal content of food*. Geneva, Switzerland. Vol.1 pp 24-35.