

UTILIZATION OF RAPESEED MEAL/CAKE IN POULTRY FEED. PART II. EFFECT OF INCORPORATING HIGHER LEVELS OF RAPESEED CAKE IN POULTRY DIET ON LAYING PERFORMANCE OF BROWN-EGG LAYER (TETRA SL)

Amal Badshah*, Aurang Zeb, Nizakat Bibi, Sajjad Ali, M A Chaudry and A Sattar

Nuclear Institute of Food and Agriculture (NIFA), Tarnab, Peshawar, Pakistan

(Received 7 May 1999; accepted 28 August 2000)

An experiment was conducted to see the effect of higher levels of rapeseed cake (15, 20 and 25%) in poultry diet on the performance of laying hen (Tetra SL). All the diets were isocaloric (within the range of 319 to 321 kcal 100 g⁻¹), isoproteinacious (17.19 to 17.31%) and the fibre contents were within the recommended range (2.98 to 3.52%). The experiment was of 28 days duration. Egg size was maximum for the 20% rapeseed cake group (62.25g) followed by the 25% group (61.24 g) and minimum for 15% rapeseed cake group (58.26 g). Egg size of standard group was 58.99 g. The egg laying percentage was 60% for standard group and 59.33% for 25% rapeseed group. No difference in birds live weight was observed due to feed variation throughout the experimental period. No mortality was recorded. Organoleptic evaluation by trained judges revealed no difference in overall acceptability of eggs by all the groups.

Key words: Rapeseed meal, Poultry diet, Egg laying.

Introduction

Controversial evidence about the nutritional value of rapeseed cake in poultry ration and its availability at a comparatively cheaper rate was the basis for conducting this experiment. A number of studies have been reported which show that rapeseed meal can be used in laying bird diets to replace a substantial quantity of soybean meal protein without any change in production performance (Hulan and Proudfoot 1980; Clandinin and Roblee 1983, Clandinin *et al* 1983; Summers 1983). There are also some reports which suggests that it may play a role in precipitating a fatty liver and/or haemorrhagic condition in layers (Pearson and Butler 1978; Papas *et al* 1979; Ibrahim *et al* 1980; Hulan and Proudfoot 1981). However, there are equal number of reports that indicate no liver problems with canola feeding (March *et al* 1975, Thomas *et al* 1978; Proudfoot *et al* 1983). Reduced egg size has also been reported with feeding of canola to laying hens (March *et al* 1975; Leeson *et al* 1978, Proudfoot *et al* 1982). From the results of our previous experiment (Badshah *et al* 1998) using brown-egg-layer, it was observed that rapeseed cake (RSC) at a level of 10% can be used in poultry ration to replace other expensive protein source with the improved production performance. Encouraged from these results, the level of rapeseed cake in the diet was increased up to 25% in the present study and the production performance was monitored and compared with that of standard. Increased levels of rapeseed cake in the poultry ration can greatly slash the cost of feed

and hence can have great economic impact on poultry production in Pakistan.

Materials and Methods

Rapeseed of commercial high-glucosinolate (60.85 μ mol g⁻¹) variety, Pakcheen, was obtained from the Mutation Breeding Section of the Nuclear Agriculture Division of Nuclear Institute for Food and Agriculture (NIFA), Peshawar. After cleaning, the seeds were expeller extracted and the material left after the oil extraction was used in the test diet formulations. A standard diet (without RSC) was also included for comparison. The compositions of the diets are given in Table 1. The chemical composition of ingredients used during the formulation of diets was taken from Scot *et al* (1982) and Malik and Chughtai (1979). A computer spread-sheet (Microsoft Excel-5) was developed for the formulation of the diets as well as for determining the calculated analysis of the feed. As depicted in Table 1, each of the diets used was isocaloric (calories 100g⁻¹ ranged between 320-321), isoproteinacious (17.19-17.31%) and their fibre content ranged between 2.98-4.35%. Assay for total glucosinolates was carried out according to the method of DeClercq and Daun (1989).

Thirty four-weeks-old hens from our previous experiment were given ten days washout period by feeding them on standard diet and then divided into four groups each replicated three times. Every group had five birds per group and were caged separately. As for the previous experiment, the size of the cage

*Author for correspondence

was 0.6 x 1.2m. The birds were maintained on natural day light condition throughout the experiment. The data given in this experiment pertains to the month of January, when the duration of the natural day light was about 11 h. Three groups of hens were fed on test diets and one group was maintained on standard ration. The RSC groups were observed for live weight (gain/loss), egg size, production and egg quality and compared with the standard group for a period of 28 days. Physical properties of eggs were determined and analysis for protein and moisture content was performed.

For the determination of egg size, average weight of all the eggs produced per group in 28 days was taken. To determine the proportion of different physical contents of eggs and protein and moisture contents, three eggs per group of the last day of the experiment were taken.

Sensory and statistical evaluation. Egg samples from RSC and standard groups were boiled, deshelled and then presented under code numbers to a panel of 9 judges trained on the subject and to whom the phenomenon of off flavour production was explained. The samples were scored on 1-10 scale for flavour, with 1 extremely disliked and 10 extremely liked (Larmond *et al* 1980). The data were statistically evaluated by the analysis of variance using co-state package and means were compared using DNMR.

Results and Discussion

The incorporation of rapeseed cake up to 25% in layer ration had no effect on live weight (gain/loss) and mortality as recorded for the subject- layer. The results (Table 2) showed maximum egg size in 20% RSC group (62.25 g) followed by 25% RSC group (61.24 g) and minimum size in 15% RSC group (58.26 g) while egg size for the standard group was 58.99 g. Egg production was recorded 60% for standard group and 59.33% RSC group. Laying performance of the other two groups were also comparable to that of the standard group. Interestingly, organoleptic quality checked through several experienced judges revealed no difference in overall acceptability of eggs from all the groups (data not given).

As for the physical properties of eggs from hens receiving different levels of RSC, a significant ($p > 0.05$) increase in the proportion of egg-white and decrease in that of egg yolk as a result of feeding on increased levels of RSC was observed. These changes seemed to be dose dependent and related to the inclusion of rapeseed cake in the diet. A decrease in moisture content (though very small) and protein content (a clear cut pattern) of egg white with the increased levels of RSC in diet (Table 3) was also noted. The moisture and protein contents of egg yolk were not influenced by RSC inclusion in ration.

Presently, the use of rapeseed meal or cake (with high glucosinolates) is subject to severe criticism in several published works. Vogt *et al* (1969 a & b) reported that with 10% rapeseed meal (glucosinolate content 0.92%) in the diet, there was a 15% reduction in the egg production, 8% lower feed conversion efficiency and 11% reduced feed intake, whereas with 20% rapeseed meal in the diet, there was 23% reduction in egg production and 26% lower feed efficiency. He also reported decreased egg quality with respect to smell and taste. The reduced performance of laying hens was attributed to the glucosinolates in the rapeseed meal. These results were quiet contradictory to our findings and to those of Summers *et al* (1971 a & b) who carried out work to see the comparative effect of feeding rapeseed meal and soybean meal on layer performance. These authors established that the poor performance of rapeseed meal-fed group was related to the amino acid imbalance in rapeseed protein rather than to some antinutrients. Although a glucosinolates threshold does not seem to have been established for layers, however in the case of broilers it was reported that depression in growth of broiler chicks fed on rapeseed meal was related to the total content of glucosinolates in the diet (Zeb *et al* 1999). It can be noted that the glucosinolate contents of all the test rations

Table 1
Composition of layer diets (%)*

| Ingredients (%) | Rapeseed cake diets | | | |
|------------------------|---------------------|--------|--------|--------|
| | Standard | 15% | 20% | 25% |
| Wheat | 38.00 | 38.00 | 38.00 | 15.00 |
| Maize | 19.00 | 15.00 | 14.00 | 37.00 |
| Rice polishing | 5.00 | 10.00 | 10.00 | 5.00 |
| Wheat bran | 5.00 | - | - | - |
| Fish meal | 4.00 | 3.00 | 1.00 | 2.00 |
| Blood meal | 3.00 | 2.00 | 1.00 | - |
| Cotton seed cake | 5.00 | - | - | - |
| Corn gluten (60%) | 4.00 | - | - | - |
| Molasses | 5.00 | 5.00 | 5.00 | 5.00 |
| Meat meal | 3.00 | 3.00 | 2.00 | 2.00 |
| Rapeseed cake | - | 15.00 | 20.00 | 25.00 |
| Dicalciumphosphate | 2.00 | 2.00 | 2.00 | 2.00 |
| Limestone | 7.00 | 7.00 | 7.00 | 7.00 |
| Total | 100.00 | 100.00 | 100.00 | 100.00 |
| Crude protein %** | 17.20 | 17.23 | 17.19 | 17.31 |
| ME (Kcal/100g) | 320.75 | 319.71 | 321.79 | 320.30 |
| Fibre % | 2.98 | 3.52 | 3.97 | 4.35 |
| Glucosinolates(###g/g) | 0.00 | 9.12 | 12.16 | 15.2 |

* To each 100 kg of diet 250g vitamineral mixture was added. The Vitamin-mixture provided per kg of the diet:- 11022 iu Vitamin A, 1100 iu vitamin D3, 11 iu vitamin E, 2.25 mg Vitamin K, 2.25 mg thiamine, 4.5 mg riboflavin, 14.25 mg pantothenic acid, 32.5 mg nicotinic acid, 4.5 mg pyridoxine, 1.25 mg folic acid, 0.25 mg biotine, 1322.5 mg choline, 55 mg Mn, 550 mg Mg, 87.5 mg Fe, 11 mg Cu, 44 mg Zn, 0.25 mg Se and 0.25 mg I.

** Calculated analysis.

Table 2

Effect of feeding different levels of rapeseed cake in the diet on production performance of brown egg layer (Tetra SL)

| Groups | Live weight record | | Production performance | |
|----------|---------------------|-------------------|------------------------|----------------|
| | Initial weight (kg) | Final weight (kg) | Egg size (g) | Production (%) |
| Standard | 1.95a | 1.98a | 58.99bc | 60.00a |
| 15% RSC | 1.98a | 1.93a | 58.26c | 58.50a |
| 20% RSC | 1.98a | 1.96a | 62.25a | 59.25a |
| 25% RSC | 1.99a | 1.95a | 61.24ab | 59.33a |

Values in the same column followed by different letters are statistically different ($p < 0.05$) from each other.

Table 3

Physical properties of eggs of brown layer receiving different levels of rapeseed cake in their diet

| Groups | Egg weight (g) | Egg white (%) | Egg yolk (%) | Egg shell (%) |
|----------|----------------|---------------|--------------|---------------|
| Standard | 60.13 | 56.99b | 29.65a | 13.09b |
| 15% RSC | 59.90 | 58.43a | 28.55b | 12.64c |
| 20% RSC | 67.00 | 57.81ab | 28.10c | 13.63a |
| 25% RSC | 63.55 | 58.49a | 27.02d | 12.02d |

Average of three determinations.

Values in the same column followed by different letters are statistically different ($p < 0.05$) from each other.

Table 4

Protein and moisture contents of egg white and yolk of brown layer receiving different levels of rapeseed cake in their diet

| Groups | Egg white | | Egg yolk | |
|----------|--------------|--------------|--------------|--------------|
| | Moisture (%) | Protein* (%) | Moisture (%) | Protein* (%) |
| Standard | 86.07a | 78.41a | 50.29b | 48.67c |
| 15% RSC | 85.49b | 77.42b | 44.08d | 59.73a |
| 20% RSC | 84.84c | 76.89c | 48.49c | 51.45b |
| 25% RSC | 84.79c | 71.44d | 53.00a | 48.21c |

*Moisture free basis.

Values in the same column followed by different letters are statistically different ($p < 0.05$) from each other.

ranged from 9 to 15 $\mu\text{g g}^{-1}$. These authors proved that the effect of glucosinolates on broiler's growth was not a linear rather a polynomeal phenomenon; where the threshold lies at 9-10 μg of glucosinolates per gram of feed. O'Neil (1957) reported that rapeseed meal could replace all of the soybean meal in a diet containing 3% animal protein without any detrimental effect on production and hatchability of eggs from

laying breeders. McGregor and Blakely (1964) and Robblee and Clandinin (1967) observed no adverse effect on the performance of turkey breeder flock using 10% rapeseed meal in their ration.

It is concluded from the experiment that RSC, having glucosinolate content in the range of those used in the present study, can be used safely upto a level of 25% without any adverse affect on the production performance of the breed used. However, further experiments are needed to confirm the results.

References

Badshah A, Zeb A, Bibi N, Ali S, Chaudry M A 1998 Utilization of rapeseed meal/cake in poultry feed.1. Effect of incorporating 10% rapeseed cake in poultry diet on the growth and laying performance of brown-egg-layer (Tetra SL). *Pak J Sci Ind Res* **41** (2) 97-101.

Clandinin D R, Robblee A R 1983 Canola meal can be good source of high quality protein for poultry. *Feedstuffs* **55** 36-37.

Clandinin D R, Robblee A R, Hardin R T, Darlington K 1983 *7th Progress Report, Research on Canola Seed Oil and Meal and Meal Fractions*. Canola Council of Canada, Winnipeg, Manitoba. Publication No. 61 pp 5-6

DeClercq D R, Daun J K 1989 Determination of the total glucosinolates content in canola by reaction with thymol and sulfuric acid. *JAOCs* **66** 788-791.

Hulan H W, Proudfoot F G 1981 Performance of laying hens fed diets containing soybean grains, rapeseed grains or rapeseed meal with and without grains. *Can J Anim Sci* **61** 1031-1040.

Hulan H W, Proudfoot F G 1980 The nutritive value of rapeseed meal for layer genotypes housed in pens. *Poultry Sci* **59** pp 585-593.

Ibrahim I K, Hodges R D, Hill R 1980 Haemorrhagic liver syndrome in laying fowl fed diets containing rapeseed meal. *Res Vet Sci* **29** 68-76.

Larmond E, Salmon R E, Klein K K 1980 Sensory evaluation of turkeys fed diets containing candle canola meal. In: *6th Progr. Rept. Res on Canola Seed, Oil, Meal and Meal Fractions*. Canola Council of Canada.

Leeson S, Slinger S J, Summers J D 1978 Utilization of whole tower rapeseed by laying hens and broiler chickens. *Can J Animal Sci* **58** 55-61.

Malik M Y, Chughtai M I D 1979 Chemical composition and nutritive value of indigenous feedstuffs. *Pak Assoc for the Advancement of Sci Lahore*.

March B E, Biely J, Soony R 1975 The effects of rapeseed meal fed during the growing and laying periods on

- mortality and egg production in chickens. *Poultry Sci* **54** 1875-1882.
- McGregor H I, Blakely R M 1964 Rapeseed oil meal as a supplement to turkey breeder ration. *Poultry Sci* **43** 189-192.
- O'Neil J B 1957 Rapeseed oil meal as a vegetable protein supplement in the diet of laying and breeding hens. *Poultry Sci* **36** 1146-1148.
- Papas A, Campbell I D, Consfield P E 1979 A study of the association of glucosinolates to rapeseed meal induced hemorrhagic liver in poultry and the influence of supplemental vitamin K. *Can J Anim Sci* **59** 133-144.
- Pearson A W, Butler E J, Curtis R F, Fenwick G R, Hobson F A, Land D G, Hall S A 1978 Effect of rapeseed meal on laying hens (*Gallus domesticus*) in relation to fatty liver hemorrhagic syndrome and egg taint. *Res Vet Sci* **25** 307-313.
- Proudfoot F G, Hulan H W, McRae K B 1982 The effect of diets supplemented with tower and/or candle rapeseed meal on performance of meat chicken breeders. *Can J Animal Sci* **62** 239-247.
- Proudfoot F G, Hulan H W, McRae K B 1983 Effect of feeding poultry diets supplemented with rapeseed meal as a primary protein source to juvenile and adult meat breeder genotypes. *Can J Anim Sci* **63** 957-965.
- Robblee A R, Clandinin D R 1967 Rapeseed meal in ration for breeder turkeys. *Can J Anim Sci* **47** 127-130.
- Scott M L, Nesheim M C, Young R J 1982 *Nutrition of the Chicken*. Publ. Scott and Associates, Ithaca, New York.
- Summers J D 1983 *Canola Meal in Layer and Broiler Diets Where Energy has not been Kept Equal. 7th Progress Rept. Research on Canola Seed Oil and Meal and Meal Fractions*. Canola Council of Canada, Winnipeg, Manitoba. Publication **61** 7.
- Summers J D, Pepper W F, Wood A S 1971 The value of rapeseed meal in broiler breeder diet. *Poultry Sci* **50** 1387-1391.
- Summers J D, Rajaratanam G, Pepper W F 1971 Evaluation of rapeseed meal as a protein supplement for laying hen diets. *Poultry Sci* **50** 1382-1386.
- Thomas D, Robblee A R, Clandinin D R 1978 Effect of low and high glucosinolate rapeseed meal on productive performance, egg quality, composition of liver and incidence of hemorrhagic liver syndrome in laying birds. *Br Poultry Sci* **19** 449-454.
- Vogt H, Schubert H J, Stute K, Rauch W 1969 Futterwert und Dinsatz von Rapsschrot in der Geflugelfutterung. 3 Mitteilung; apsschrot in der Legehennenfütterung. *Arch Geflugelk* **33** 119-124.
- Vogt H, Schubert H J, Stute K 1969 Futterwert and Einsatz von Rapsschrot in der Geflugelfütterung. 4 Mitteilung; Zweiter Legehennen Versuch mit Rapsschrot. *Arch Geflugelk* **33** 392-395.
- Zeb A, Sattar A, Ter Meulen U 1999 Effect of feeding different levels of rapeseed meal on the performance of broiler chicks. *Arch fue Geflugelkd* **63** (2) 1-5. ISBN 003-9098.