

CONDENSED BENZOPYRANS VI. 8H-PYRANO[3,2-f]QUINOXALINES

Misbahul Ain Khan^{ac*} and Edvaldo Cavalcanti Caldas^b

^aSecção de Química, Instituto Militar de Engenharia, Praia Vermelha, Rio de Janeiro, RJ, Brasil. CEP. 22 290

^bUnigranrio -Universidade do Grande Rio, Departamento de Química, Rua Prof. Jose de Souza Herdy 1160, 25 de Agosto, Duque de Caxias, RJ, Brasil, CEP 25071-200

^c459-A, Block C, Faisal Town, Lahore, Pakistan

(Received 12 August 2000; accepted 10 April 2001)

The basic ring system 8H-pyrano[3,2-f]quinoxaline and its alkyl, aryl and oxo derivatives (5a-5d) were prepared by the reaction of 5,6-diaminocoumarin with appropriate 1,2-dicarbonyl compounds.

Key words: Pyrano[3,2-f]quinoxalines, coumarins, ¹H NMR Spectra.

Introduction

Of the various isomeric pyranoquinoxalines only a handful have been mentioned in literature. A 3,4-dihydro-3-methyl-1-ol, derivative of pyrano [3,4-b]quinoxaline (**1**), was obtained and spectroscopically identified from the action of pyransome dehydratase, of *Phanerochaete chrysosporium* on osones and the labile intermediate trapped by *o*-phenylenediamine (Gabriel *et al* 1993). Some derivatives of the system pyrano [3,2-f] quinoxaline (**2**) and pyrano [2,3-f] quinoxaline (**3**) were prepared from the 6- or 5-hydroxyquinoxalines respectively by their reactions with α,β -unsaturated aldehydes and organotitanium compounds such as titanium tetraethoxide. These pyranoquinoxalines, however, contained aryl groups at 8-(or 9-) position ($R_1=R_2=aryl$), and various such derivatives are claimed to display photochromism and to have use in ophthalmic optics (Guglielmetti *et al* 1993; Pozzo *et al* 1994; Hara and Momota 1996; Samat *et al* 1997; Pozzo *et al* 1997).

During our continuous interest in condensed benzopyrans containing nitrogen heterocyclic systems we had earlier reported the synthesis of systems with fused pyrrole (Khan and de B Morley 1978, 1979), pyridine (Khan and Gemal 1977, 1978), imidazole (Khan and Caldas 1987) and 1,2,3-triazole rings (Khan and Caldas 1986). In this communication we wish to report our results on the synthesis of 8H-pyrano[3,2-f]quinoxaline ring system.

Experimental

The ¹H nuclear magnetic resonance spectra were obtained on a Hitachi Perkin-Elmer R20B spectrometer operating at 60 MHz with tetramethylsilane as internal standard. The infrared

absorption spectra were taken by Perkin-Elmer model 727 spectrophotometer. The samples were measured in potassium bromide disks. Melting points were determined with a Fisher-Johns apparatus and are uncorrected. Elemental analyses were carried on a Perkin-Elmer model 240.

8-Oxo-8H-Pyrano[3,2-f]quinoxaline (5a): A mixture of 0.2 g of 5,6-diaminocoumarin (**4**) (Khan and Caldas 1986), 0.66 g of glyoxal, 10 ml of water and a few drops of hydrochloric acid was heated under reflux for two hours. On cooling the precipitate was filtered off and crystallized from ethanol (activated charcoal). **5a** was obtained as colorless crystals, mp 215-216°C. Yield 0.05 g (20%). Infra red (cm⁻¹): 1760 (CO lactone); ¹H nmr (deuteriochloroform): δ 6.58 (d, 1H, J=10Hz, H-9), 7.68 (d, 1H, J=10Hz, H-6), 8.16 (d, 1H, J=10Hz, H-5), 8.84 (s, 2H, H-2 & H-3), 8.85 (d, 1H, J=10 Hz, H-10).

Anal. Calcd. for C₁₁H₆N₂O₂: C, 66.67, H, 3.05, N, 14.14; Found, C, 66.54, H, 3.07, N, 13.91.

2,3-Dimethyl-8-oxo-8H-pyrano[3,2-f] quinoxaline (5b): Using the method for the preparation of **5a**, from 0.1g of **4** and 0.5 ml of diacetyl, 0.1 g (80%) of **5b**, was obtained as yellow crystals from dimethylsulfoxide. mp 225-226°C. Infra-red (cm⁻¹): 1720 (CO lactone), ¹H nmr (trifluoroacetic acid): δ 2.70 (s, 3H, CH₃), 2.75 (s, 3H, CH₃), 6.60 (d, 1H, J=10 Hz, H-9), 7.68 (d, 1H, JH=9Hz, H-6), 8.05 (d, 1H, J=9 Hz, H-5), 8.82 (d, 1H, J=10 Hz, H-10).

Anal. Calcd. for C₁₃H₁₀N₂O₂: C, 69.02, H, 4.46, N, 12.38; Found, C, 68.87, H, 4.45, N, 12.03.

2,3-Diphenyl-8-oxo-8H-pyrano[3,2-f]quinoxaline (5c): Similarly from 0.10g of **4** and 0.12 g of benzil, 0.12 g of **5c** was obtained as yellow crystals from dimethylsulfoxide, mp 293-294°C, yield 60%. Replacing water with ethanol for the

*Author for correspondence, ^cPresent address

reaction raised the yield of 5c to 73%. Infrared (cm^{-1}): 1740 (CO lactone); ^1H nmr (trifluoroacetic acid): δ 6.59 (d, 1H, $J=10$ Hz, H-9), 7.20 (m, 10H, Ph), 7.72 (d, 1H, $J=10$ Hz, H-6), 8.15 (d, 1H, $J=10$ Hz, H-5), 8.85 (d, 1H, $J=10$ Hz, H-10).

Anal. Calcd. for $\text{C}_{23}\text{H}_{14}\text{N}_2\text{O}_2$: C, 78.84, H, 4.03, N, 8.00; Found, C, 78.66, H, 4.09, N, 7.93.


8H-Pyrano[3,2-f]quinoxaline-2,3,8-trione-(5d):

Following the method for the synthesis for 5a-5c, 0.1 g of 4 and 0.5 mL of diethyl oxalate, 5d was obtained as rose colored crystals from dimethylsulfoxide, mp >300°C, yield 84%. Infrared (cm^{-1}): 3100-3000 (br. NH) and 1750-1650 (br. CO amide & lactone). Not very soluble in usual organic solvents for ^1H nmr spectrum.

Anal. Calcd. for $\text{C}_{11}\text{H}_6\text{N}_2\text{O}_4$: C, 57.40, H, 2.63, N, 12.17; Found, C, 57.62, H, 2.70, N, 11.90.

Results and Discussion

The key starting material for the synthesis of various 8H-pyran[3,2-f]quinoxalines is 5,6-diamino coumarin (4) which was earlier obtained from coumarin by its successive nitration and reduction (Khan and Caldas 1986). This diaminocoumarin was condensed with 1,2-dicarbonyl compounds for the construction of the quinoxaline ring (Cheeseman 1963; Cheeseman and Werstiuk 1978). The synthesis of various 8H-pyran[3,2-f]quinoxalines is presented in the scheme 1.



Scheme 1

The condensation of 4 with glyoxal produced the basic ring system of 8H-pyran[3,2-f]quinoxalines (5a). Its infrared spectrum was devoid of any absorption bands in the 3200 - 3500 cm^{-1} region indicating disappearance of the two amino groups. This evidence together with its ^1H nuclear magnetic resonance spectrum which displayed four doublets at δ 6.58, 8.85, 8.16 and 7.86, with coupling constants of 10 Hz for the protons at positions 9, 10, 5 and 6 respectively and two protons singlet at 8.84 for the protons at 2 and 3 positions, indicated the formation of the desired ring system 5a. The infrared and ^1H nuclear magnetic resonance spectra for the compounds 5b-5d, together with their elemental analyses helped, in establishing their structure (see experimental). The compound 5d exists mainly in the "dilactam" form (infrared absorption in the region 3100-3000 cm^{-1} and 1750-1650 cm^{-1} for "lactam" NH and CO respectively).

Acknowledgements

Continuous support from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) of Brazil is gratefully acknowledged. Edvaldo Cavalcanti Caldas also thanks CNPq for the fellowship.

References

- Cheeseman GWH 1963 Recent advances in quinoxaline chemistry *Adv Heterocyclic Chem* **2** 203.
- Cheeseman GWH, Werstiuk ESG 1978 Quinoxaline chemistry development 1963-1978. *Adv Heterocyclic Chem* **22** 367.
- Gabriel J, Vole J, Sedmera P, Daniel G, Kubatova E 1993 Pyranosone dehydratase from the basidiomycete *Phanerochaete chrysosporium*: Improved purification and identification of 6-Deoxy-D-glucosone and D-xylosone reaction products. *Arch Microbiol* **160** 27-34.
- Guglielmetti R, Pozzo J L, Samat A 1993 Preparation of photochromic heterocyclic-fused chromenes and their use in ophthalmic optics. *Eur Pat Appl. EP 562,915 Chem. Abstr* **120** 106977 (1994).
- Hara T, Momota J 1996 Preparation of 2-Phenyl-2-(2-naphthyl) naphtha [1,2-b] pyran Derivatives and their heterocyclic analogs as photochromic materials. *Jpn Kokai Tokkyo Koho JP 08, 295, 690 Chem. Abstr* **126** 89395 (1997).
- Khan MA, Caldas EC 1986 1H-, 7H-Pyran[3,2-e]-V-benzotriazol-7-one, A new ring system. *Bull Korean Chem Soc* **7** 412-413
- Khan MA, Caldas EC 1987 Condensed benzopyrans V. Synthesis of some derivatives of 7H-pyran[3,2-e] benzimidazol-7-one. A new ring system. *Chem. Scr* **27** 269-270.

- Khan MA, DeB Morley ML 1978 Condensed benzopyrans III. 3*H*-(1)-benzopyrano-[3,4-*b*] pyrrol-4-ones. *J Heterocyclic Chem* **15** 1399-1401.
- Khan MA, DeB Morley ML 1979 Condensed benzopyrans IV. Synthesis of some derivatives of 7*H*-, and 9*H*-Pyrano [3,2-*e*] indoles. *J Heterocyclic Chem* **16** 997-999.
- Khan MA, Gemal AL 1977 Synthesis of 5*H*-benzopyrano [3,4-*b*] pyridin-5-one and its derivatives. *J Heterocyclic Chem* **14** 1009-1010.
- Khan MA, Gemal AL 1978 Condensed benzopyrans II. Some derivatives of 3*H*-pyrano[3,2-*b*] quinoline. *J Heterocyclic Chem* **15** 159-160.
- Pozzo JL, Lokshin V A, Guglielmetti R 1994 New photochromic 2,2-diphenyl-[2*H*]-chromenes annealed with nitrogenated six-membered ring. *mol. Cryst Liq Cryst Sci Technol Sect A* **246** 75-8. *Chem Abstr* **122** 92549 (1995).
- Pozzo JL, Samat A, Guglielmetti R, Dubest R, Aubard J 1997 Synthesis and photochromic behaviour of naphthopyrans, pyranoquinolines, and pyrano-quinoxalines *Helv Chim Acta* **80** 725-738.
- Samat A, Garros G, Pommier H, Pottiker E, Pepe G, Guglielmetti R, Rajzmann M 1997. Color prevision of activated forms of photochromic spirooxazines and chromenes. *mol. Cryst Liq Cryst Sci Technol Sect A* **298** 297-304 *Chem Abstr* **127** 227205 (1997).