

## KINETIC STUDY OF REDUCTION OF BIS(PENTANE-2,4-DIONATO) DIAQUO MANGANESE (III) COMPLEX WITH SODIUM THIOSULPHATE

M Akhyar Farrukh\* and Iftikhar Imam Naqvi

Department of Chemistry, University of Karachi, Karachi-75270, Pakistan

(Received 02 December 2000; accepted 6 August 2001)

Kinetics of several oxidation-reduction reactions pertaining to manganese have been reported but very few of them involve manganese (III) in perchloric acid. This may be because free manganic ions can not exist in significant concentration in aqueous solutions. Manganese (III) forms a number of complexes with fluoride, chloride, phosphate, pyrophosphate, sulphate, porphyrin, heteropoly anions (Baker 1961; Brauer 1963; Kemmit 1973)

We have now investigated the kinetics and mechanisms of the reduction of bis (pentane 2,4-dionato) diaquo manganese (III) complex by sodium thiosulphate. Kinetic investigation of the electron transfer reactions of metal complexes of  $\beta$ -diketones are sparse and this presentation is one of the few studies of such reactions using simple reductant.

The tris (pentane-2,4-dionato) manganese (III) complex  $[\text{Mn}(\text{pd})_3]$  was prepared as previously described (Charles and Bryant 1963). Bis (pentane 2,4-dionato) diaquo manganese (III) complex  $[\text{Mn}(\text{pd})_2(\text{H}_2\text{O})_2]^+$  was prepared from tris (pentane-2,4-dionato) manganese (III) complex by substitution reaction method reported earlier (Cartledge 1952).

Molar absorptivity value of reduced form of complex was determined to convert the absorbance rate data to the concentration rate data. It was found to be  $90.2 \text{ lit mol}^{-1} \text{ cm}^{-1}$  at  $\lambda_{\text{max}} = 298.5 \text{ nm}$

Absorption spectra were recorded at 298.5nm by using Shimadzu UV-160, UV-visible spectrophotometer.

Different combinations of both complex and reductant (sodium thiosulphate) were prepared by taking constant concentration of complex and varying the concentration of the reductant. Perchloric acid was used as a source of hydrogen ion for the pH range of 2-2.5 and all solutions were adjusted to an ionic strength of  $0.5 \text{ mole dm}^{-3}$  by the addition of  $\text{NaClO}_4$ . The absorbance of each set of reaction mixture was recorded at 298.5nm. Mn (III) was analyzed potentiometrically. In this technique the complex  $[\text{Mn}(\text{pd})_3]$  was reduced by L-ascorbic acid and potential differences were read with the aid of a potentiometer (Kent EIL 7020). Analysis of the above complex gave Mn 17.1%, where as calculated amount is 15.6%.

Data were collected using initial rate method by plotting the

**Table 1**  
Kinetic rate data between the reaction of  $\text{S}_2\text{O}_3^{2-}$  with  $\text{Mn}(\text{pd})_2(\text{H}_2\text{O})_2^+$ ,  $\text{NaClO}_4=0.5\text{M}$

| S.No | Composition of reaction mixture (ml) |         |              | [Complex] M<br>$10^4$ | [Thiosulphate] M<br>$10^4$ | Initial rate<br>A $\text{Sec}^{-1}$ | $k' \text{ Sec}^{-1}$<br>Calculated                              | $k' \text{ Sec}^{-1}$<br>Graphical |
|------|--------------------------------------|---------|--------------|-----------------------|----------------------------|-------------------------------------|--|------------------------------------|
|      | $\text{NaClO}_4$                     | Complex | Thiosulphate |                       |                            |                                     |  |                                    |
| A    | 2.3                                  | 0.5     | 0.2          | 8.33                  | 6.66                       | 1.083                               | Rate is independent of the initial concentration of thiosulphate |                                    |
| B    | 2.2                                  | 0.5     | 0.3          | 8.33                  | 10                         | 1.083                               |  |                                    |
| C    | 2.1                                  | 0.5     | 0.4          | 8.33                  | 13.3                       | 1.089                               |  |                                    |
| D    | 2.0                                  | 0.5     | 0.5          | 8.33                  | 16.6                       | 1.08                                | 14.37  |                                    |
| E    | 2.4                                  | 0.1     | 0.5          | 1.66                  | 16.6                       | 0.244                               | 16.29  | 13.3                               |
| F    | 1.8                                  | 0.7     | 0.5          | 11.66                 | 16.6                       | 1.40                                | 13.31  |                                    |
| G    | 1.6                                  | 0.9     | 0.5          | 15.0                  | 16.6                       | 1.749                               | 12.92  |                                    |

$\lambda_{\text{max}} = 298.5 \text{ nm}$ ; Complex = 0.005M;  $\text{S}_2\text{O}_3^{2-} = 0.01\text{M}$ ; pH= 2-2.5;  $\epsilon = 90.2 \text{ lit mol}^{-1} \text{ cm}^{-1}$

experiments (A-D) with constant complex, [bis(Pentane-2,4-dionato) diaquo manganese (III)] and variable reductant (sodium thiosulphate) concentrations and another four experi-

\*Authors for correspondence

ments (D-G) with constant reductant and variable complex concentration, lead to the results reported here in.

In the first case (experiments A-D) with constant complex concentration, the plot shows that there is no change in the




Fig 1(a). of Table 1. Plot of rate Vs concentration at constant complex and variable reductant concentration




Fig 1 (b). of Table 1. Plot of rate Vs concentration at constant reductant and variable complex concentration




Fig 2(a). of Table 2. Plot of pseudo- first order rate constant Vs reductant concentration at constant complex concentration




Fig 2(b). of Table 2. Plot of pseudo- first order rate constant Vs ln [reductant] at constant complex concentration

Table 2

Indirectly determination of order of reaction of reductant at constant complex concentration

| S.No | [Complex]M<br>10 <sup>4</sup> | [Thiosulphate]M<br>10 <sup>4</sup> | k' Sec <sup>-1</sup><br>Graphical | ln k' | ln [thiosulphate] |
|------|-------------------------------|------------------------------------|-----------------------------------|-------|-------------------|
| 1    | 8.33                          | 16.66                              | 13.3                              | 2.59  | 2.81              |
| 2    | 8.33                          | 11.66                              | 8.9                               | 2.2   | 2.46              |
| 3    | 8.33                          | 25                                 | 18.5                              | 2.91  | 3.22              |


rate when the concentration of the reductant is varied. So, the order of reaction with respect to reductant is zero. As shown in Table 1 and in Fig 1(a). In the second case (experiments D-G), rate (absorbance persecond) increases in a linear manner with the concentration of the complex as shown in Table 1 and in Fig 1(b). This shows that order of reaction with respect to complex is first order and that the overall order of reaction is pseudo-first order.

When the pseudo-first order rate constants, k', determined graphically, given in Table 2, of previous different set of experiments are plotted versus the reductant concentrations, as shown in

Fig 2(a), a straight line is obtained passing through the origin. The straight line indicates that the order of reaction with respect to reductant is also first order. The slope of this later plot gives the overall second order reaction.

Another approach to determining the order of reaction of the reductant is a  $\ln k' - \ln$  plot as shown in Fig. 2(b). The slope of the plot of  $\ln k'$  versus  $\ln$  [reductant] should equal to 'n' order of reaction. The value of slope determined by this method is equal to '1' so this is again an indication of first order reaction with respect to reductant (sodium thiosulphate) and over all order of reaction is second order. The intercept gives the value of second order rate constant, amounting to  $7.5 \times 10^3 \text{ M}^{-1}\text{S}^{-1}$

In view of the above mentioned experimental results we propose the following mechanism for the reaction.



**References**

- Baker LCW 1961 *Advances in Chemistry of the Coordination Compound*. Macmillan, New York, pp 604-612.
- Brauer G 1963 *Handbook of Preparative Inorganic Chemistry*. Academic Press, **1 & 2** 2<sup>nd</sup> ed.
- Cartledge G H 1952 Bis(pentaenedione)diaquo manganese (III) complex. *J Am Chem Soc* **74** 6015-6018.
- Charles R C, Bryant B E 1963 Acetylacetonatomanganese (III) *Inorg Synth* **7** 183-184.
- Kemitt RD 1973 *Comprehensive Inorganic Chemistry*. Pergamon Press, **3** pp 786 and 797.